


AMENDMENTS TO THE CLAIMS

The following listing of claims will replace all prior versions and listings of claims in the application.

LISTING OF CLAIMS


1. (Currently Amended) A compound of the formula (I)









I

or a pharmaceutically acceptable salt, prodrug or ester thereof, wherein:

G is selected from the group consisting of



wherein Q is NH or O and Q' is H, C₁-C₆ alkyl, phenyl, or phenyl-C₁-C₄-alkyl;



wherein R' and R'' are independently H or C₁-C₄ alkyl;

B is (CH₂)_m(CH=CH)_pY, wherein m = 1,2,3, p = 0,1, Y is CH₂ or CO[.] ;

X is CH₂ or C=O;

R₁ is selected from the group consisting of H, C₁-C₄ alkyl, C₁-C₄ alkoxy, OH, halogen, and CF₃;

A is CH₂, NH, O, or S(O)_n wherein n is zero, 1 or 2[.] ;


R₂ is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, aryl or C₅-C₇ monocyclic

heteroaryl ring containing one to three heteroatoms selected from O, S, and N, unsubstituted or optionally substituted by one to three substituents independently selected from the group consisting of H, C₁-C₄ alkyl, C₁-C₄ alkoxy, OH, halogen, and CF₃;







R is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₄ alkynyl, aryl or aryl-C₁-C₄ alkyl.

2. (Currently Amended) A compound according to claim 1, wherein

G is selected from the group consisting of



wherein Q is NH or O and Q' is selected from the group consisting of H, C₁-C₆ alkyl, phenyl, and phenyl-C₁-C₄-alkyl;



wherein R' is as defined in claim 1;


B is (CH₂)_q where q is 2,3,4;

X, R₁, A and R are as defined [[above]] in claim 1;

R₂ is a phenyl or pyridine ring unsubstituted or optionally substituted as defined in claim 1.

3. (Previously Presented) A compound according to claim 2, wherein

G is selected from the group consisting of



wherein B, X, R₁, A and R are as defined in claim 2.

4. (Previously Presented) The compound as recited in claim 1 wherein the compound is selected from the group consisting of:

3-phenyl-N-{4-[2-(2-pyridinylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

3-phenyl-N-{4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

3-phenyl-N-{4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

N-{4-[2-(1H-imidazol-2-ylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine

N-{4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine

N-{4-[2-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine

3-(3-pyridinyl)-N-{4-[2-(2-pyridinylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

3-(3-pyridinyl)-N-{4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

3-(3-pyridinyl)-N-{4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

N-{4-[2-(1H-imidazol-2-ylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine

N-{4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine

N-{4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine

N-{3-oxo-4-[2-(2-pyridinylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine


N-{3-oxo-4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine

N-{3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-

phenyl-beta-alanine
 N-{4-[2-(1H-imidazol-2-ylamino)ethyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-
 3-phenyl-beta-alanine
 N-{4-[3-(1H-imidazol-2-ylamino)propyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-
 yl}-3-phenyl-beta-alanine
 N-{4-[4-(1H-imidazol-2-ylamino)butyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-
 3-phenyl-beta-alanine
 N-{3-oxo-4-[2-(2-pyridinylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-
 pyridinyl)-beta-alanine
 N-{3-oxo-4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-
 (3-pyridinyl)-beta-alanine
 N-{3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-
 pyridinyl)-beta-alanine
 N-{4-[2-(1H-imidazol-2-ylamino)ethyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-
 3-(3-pyridinyl)-beta-alanine
 N-{4-[3-(1H-imidazol-2-ylamino)propyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-
 yl}-3-(3-pyridinyl)-beta-alanine
 N-{4-[4-(1H-imidazol-2-ylamino)butyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-
 3-(3-pyridinyl)-beta-alanine
 3-({3-oxo-4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-
 yl}oxy)-3-phenylpropanoic acid;
 3-({3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-
 3-phenylpropanoic acid;
 3-({3-oxo-4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-
 yl}oxy)-3-phenylpropanoic acid;
 3-({3-oxo-4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-
 3-(3-pyridinyl)propanoic acid;
 3-({3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-
 3-(3-pyridinyl)propanoic acid;
 3-({3-oxo-4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-
 yl}oxy)-3-(3-pyridinyl)propanoic acid;


3-({3-oxo-4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-(pyridinyl)propanoic acid;
 3-({3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-phenylpropanoic acid;
 3-({3-oxo-4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-phenylpropanoic acid;
 3-({3-oxo-4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-phenylpropanoic acid;
 3-({3-oxo-4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-(3-pyridinyl)propanoic acid;
 3-({3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-(3-pyridinyl)propanoic acid;
 3-({3-oxo-4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-(3-pyridinyl)propanoic acid;
 3-({3-oxo-4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-(3-pyridinyl)propanoic acid;
 4-{3-oxo-4-[4-(pyridin-2-ylamino)-butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenylbutanoic acid;
 4-{3-oxo-4-[3-(1H-imidazol-2-ylamino)-propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenylbutanoic acid;
 4-{3-oxo-4-[4-(1H-imidazol-2-ylamino)-butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenylbutanoic acid;
 4-{3-oxo-4-[4-(pyridin-2-ylamino)-butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)butanoic acid;
 4-{3-oxo-4-[3-(1H-imidazol-2-ylamino)-propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)butanoic acid; and
 4-{3-oxo-4-[4-(1H-imidazol-2-ylamino)-butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)butanoic acid;
 either as a single isomer or as a mixture thereof, or a pharmaceutically acceptable salt thereof.

5. (Currently Amended) A pharmaceutical composition comprising a therapeutically effective amount of a compound or a pharmaceutically acceptable salt, prodrug or ester thereof having the formula (I):









wherein:

G is selected from the group consisting of



wherein Q is NH or O and Q' is H, C₁-C₆ alkyl, phenyl, or phenyl-C₁-C₄-alkyl;



wherein R' and R'' are independently H or C₁-C₄ alkyl;

B is (CH₂)_m(CH=CH)_pY, wherein m = 1,2,3, p = 0,1, Y is CH₂ or CO[.];

X is CH₂ or C=O;

R₁ is selected from the group consisting of H, C₁-C₄ alkyl, C₁-C₄ alkoxy, OH, halogen, and CF₃;

A is CH₂, NH, O, S(O)_n wherein n is zero, 1 or 2[.];


R₂ is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, aryl or C₅-C₇ monocyclic heteroaryl ring containing one to three heteroatoms selected from O, S, and N, unsubstituted or optionally substituted by one to three substituents independently selected from the group consisting of H, C₁-C₄ alkyl, C₁-C₄ alkoxy, OH, halogen,

and CF₃;






R is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₄ alkynyl, aryl or aryl-C₁-C₄ alkyl.

6. (Previously Presented) A pharmaceutical composition of claim 5 wherein:


G is selected from the group consisting of



wherein Q is NH or O and Q' is selected from the group consisting of H, C₁-C₆ alkyl, phenyl, and phenyl-C₁-C₄-alkyl;



and




B is (CH₂)_q where q is 2,3,4;


R₂ is a phenyl or pyridine ring unsubstituted or optionally substituted by one to three substituents independently selected from the group consisting of H, C₁-C₄ alkyl, C₁-C₄ alkoxy, OH, halogen, and CF₃.

7. (Previously Presented) A pharmaceutical composition of claim 6 wherein:

G is selected from the group consisting of



and



8. (Previously Presented) A pharmaceutical composition comprising a therapeutically effective amount of a compound or a pharmaceutically acceptable salt, prodrug or ester thereof as recited in claim 5 wherein the compound is selected from the group consisting of

3-phenyl-N-{4-[2-(2-pyridinylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

3-phenyl-N-{4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

3-phenyl-N-{4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

N-{4-[2-(1H-imidazol-2-ylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine

N-{4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine

N-{4-[2-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine

3-(3-pyridinyl)-N-{4-[2-(2-pyridinylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

3-(3-pyridinyl)-N-{4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

3-(3-pyridinyl)-N-{4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

N-{4-[2-(1H-imidazol-2-ylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine

N-{4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine

N-{4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine

N-{3-oxo-4-[2-(2-pyridinylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine

N-{3-oxo-4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine

N-{3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine

N-{4-[2-(1H-imidazol-2-ylamino)ethyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-

3-phenyl-beta-alanine

N-{4-[3-(1H-imidazol-2-ylamino)propyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine

N-{4-[4-(1H-imidazol-2-ylamino)butyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine

N-{3-oxo-4-[2-(2-pyridinylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine

N-{3-oxo-4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine

N-{3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine

N-{4-[2-(1H-imidazol-2-ylamino)ethyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine

N-{4-[3-(1H-imidazol-2-ylamino)propyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine

N-{4-[4-(1H-imidazol-2-ylamino)butyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine

3-({3-oxo-4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-phenylpropanoic acid;

3-({3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-phenylpropanoic acid;

3-({3-oxo-4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-phenylpropanoic acid;

3-({3-oxo-4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-(3-pyridinyl)propanoic acid;

3-({3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-(3-pyridinyl)propanoic acid;


3-({3-oxo-4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-(3-pyridinyl)propanoic acid;

3-({3-oxo-4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-(pyridinyl)propanoic acid;

3-({3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-phenylpropanoic acid;
 3-({3-oxo-4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-phenylpropanoic acid;
 3-({3-oxo-4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-phenylpropanoic acid;
 3-({3-oxo-4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-(3-ppyrindinyl)propanoic acid;
 3-({3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-(3-pyridinyl)propanoic acid;
 3-({3-oxo-4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-(3-pyridinyl)propanoic acid;
 3-({3-oxo-4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-(3-pyridinyl)propanoic acid;
 4-{3-oxo-4-[4-(pyridin-2-ylamino)-butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenylbutanoic acid;
 4-{3-oxo-4-[3-(1H-imidazol-2-ylamino)-propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenylbutanoic acid;
 4-{3-oxo-4-[4-(1H-imidazol-2-ylamino)-butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenylbutanoic acid;
 4-{3-oxo-4-[4-(pyridin-2-ylamino)-butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)butanoic acid;
 4-{3-oxo-4-[3-(1H-imidazol-2-ylamino)-propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)butanoic acid; and
 4-{3-oxo-4-[4-(1H-imidazol-2-ylamino)-butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)butanoic acid;
 either as a single isomer or as a mixture thereof.


9. (Currently Amended) A method for treating a condition mediated by the $\alpha_v\beta_3$ integrin in a mammal in need of such treatment, including a human, comprising administering to said mammal an effective $\alpha_v\beta_3$ inhibiting amount of a compound

of formula (I)









wherein:

G is selected from the group consisting of:



wherein Q is NH or O and Q' is H, C₁-C₆ alkyl, phenyl, or phenyl-C₁-C₄-alkyl;



wherein R' and R'' are independently H or C₁-C₄ alkyl;

B is (CH₂)_m(CH=CH)_pY, wherein m = 1, 2, 3, p = 0, 1, Y is CH₂ or CO[.];

X is CH₂ or C=O;

R₁ is selected from the group consisting of H, C₁-C₄ alkyl, C₁-C₄ alkoxy, OH, halogen, and CF₃;


A is CH₂, NH, O, S(O)_n wherein n is zero, 1 or 2[.];

R₂ is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, aryl or C₅-C₇ monocyclic heteroaryl ring containing one to three heteroatoms selected from O, S, and N, unsubstituted or optionally substituted by one to three substituents selected independently from H, C₁-C₄ alkyl, C₁-C₄ alkoxy, OH, halogen, and CF₃;







R is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₄ alkynyl, aryl and aryl-C₁-C₄ alkyl.

10. (Previously Presented) The method of claim 9 wherein :

G is selected from the group consisting of



wherein Q is NH or O and Q' is selected from the group consisting of H, C₁-C₆ alkyl, phenyl, and phenyl-C₁-C₄-alkyl;




B is (CH₂)_q where q is 2,3,4

R₂ is a phenyl or pyridine ring unsubstituted or optionally substituted by one to three substituents independently selected from the group consisting of H, C₁-C₄ alkyl, C₁-C₄ alkoxy, OH, halogen, and CF₃.

11. (Previously Presented) The method of claim 9 wherein :

G is selected from the group consisting of



12. (Previously Presented) The method according to claim 9 wherein the compound is selected from the group consisting of

3-phenyl-N-{4-[2-(2-pyridinylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

3-phenyl-N-{4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

3-phenyl-N-{4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine

N-{4-[2-(1H-imidazol-2-ylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-

phenyl-beta-alanine
 N-{4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine
 N-{4-[2-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine
 3-(3-pyridinyl)-N-{4-[2-(2-pyridinylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine
 3-(3-pyridinyl)-N-{4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine
 3-(3-pyridinyl)-N-{4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-beta-alanine
 N-{4-[2-(1H-imidazol-2-ylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine
 N-{4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine
 N-{4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine
 N-{3-oxo-4-[2-(2-pyridinylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine
 N-{3-oxo-4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine
 N-{3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine
 N-{4-[2-(1H-imidazol-2-ylamino)ethyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine
 N-{4-[3-(1H-imidazol-2-ylamino)propyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine
 N-{4-[4-(1H-imidazol-2-ylamino)butyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenyl-beta-alanine
 N-{3-oxo-4-[2-(2-pyridinylamino)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine

N-{3-oxo-4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine
 N-{3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine
 N-{4-[2-(1H-imidazol-2-ylamino)ethyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine
 N-{4-[3-(1H-imidazol-2-ylamino)propyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine
 N-{4-[4-(1H-imidazol-2-ylamino)butyl]-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)-beta-alanine
 3-({3-oxo-4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-phenylpropanoic acid;
 3-({3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-phenylpropanoic acid;
 3-({3-oxo-4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-phenylpropanoic acid;
 3-({3-oxo-4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-(3-pyridinyl)propanoic acid;
 3-({3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-(3-pyridinyl)propanoic acid;
 3-({3-oxo-4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-(3-pyridinyl)propanoic acid;
 3-({3-oxo-4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}oxy)-3-(pyridinyl)propanoic acid;
 3-({3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-phenylpropanoic acid;
 3-({3-oxo-4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-phenylpropanoic acid;
 3-({3-oxo-4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-phenylpropanoic acid;
 3-({3-oxo-4-[3-(2-pyridinylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-

yl}sulfanyl)-3-(3-ppyridinyl)propanoic acid;
 3-({3-oxo-4-[4-(2-pyridinylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-(3-pyridinyl)propanoic acid;
 3-({3-oxo-4-[3-(1H-imidazol-2-ylamino)propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-(3-pyridinyl)propanoic acid;
 3-({3-oxo-4-[4-(1H-imidazol-2-ylamino)butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}sulfanyl)-3-(3-pyridinyl)propanoic acid;
 4-{3-oxo-4-[4-(pyridin-2-ylamino)-butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenylbutanoic acid;
 4-{3-oxo-4-[3-(1H-imidazol-2-ylamino)-propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenylbutanoic acid;
 4-{3-oxo-4-[4-(1H-imidazol-2-ylamino)-butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-phenylbutanoic acid;
 4-{3-oxo-4-[4-(pyridin-2-ylamino)-butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)butanoic acid;
 4-{3-oxo-4-[3-(1H-imidazol-2-ylamino)-propyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)butanoic acid; and
 4-{3-oxo-4-[4-(1H-imidazol-2-ylamino)-butyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}-3-(3-pyridinyl)butanoic acid;
 either as a single isomer or as a mixture thereof, or a pharmaceutically acceptable salt thereof.

13. (Previously Presented) The method according to claim 9, wherein the condition treated is bone resorption, osteoporosis, humoral hypercalcemia of malignancy, Paget's disease, tumor metastasis, neoplasia (solid tumor growth), angiogenesis including tumor angiogenesis, diabetic retinopathy, arthritis, psoriasis and periodontal disease, or smooth muscle cell migration.

14. (Previously Presented) The method according to claim 12, wherein the condition treated is bone resorption, osteoporosis, humoral hypercalcemia of malignancy, Paget's disease, tumor metastasis, neoplasia (solid tumor growth),

angiogenesis, diabetic retinopathy, arthritis, psoriasis and periodontal disease, or smooth muscle cell migration.

15. (Previously Presented) A combined method of treatment of cancer or of controlling the growth of a neoplasm in a mammal suffering from cancer, said method comprising administering simultaneous, separately or sequentially,

- 1) a compound of formula (I) as defined in claim 1 or a pharmaceutically acceptable salts thereof and
- 2) an additional antitumor agent; in amounts and close enough together in time sufficient to produce a therapeutically useful effect.

16. (Original) The method according to claim 15, wherein the additional antitumor agent is selected from the group consisting of an antineoplastic topoisomerase II inhibitor, an antineoplastic antimicrotubule agent, an antineoplastic alkylating agent, an antineoplastic antimetabolite and an antineoplastic topoisomerase I inhibitor.

17. (Original) A product containing a compound of formula (I) as defined in claim 1 or a pharmaceutically acceptable salt thereof, and an effective antineoplastic amount of additional antitumor agent as a combined preparation for simultaneous, separate or sequential use in anti-cancer therapy.

18. (Original) The product according to claim 17, wherein the additional antitumor agent is selected from an antineoplastic topoisomerase II inhibitor, an antineoplastic antimicrotubule agent, an antineoplastic alkylating agent, an antineoplastic antimetabolite and an antineoplastic topoisomerase I inhibitor.

19. (Previously Presented) The method according to claim 9, wherein the condition treated is tumor angiogenesis.